

Correlation between mutation pressure, selection pressure and occurrence of amino acids

Aleksandra Nowicka¹, Paweł Mackiewicz¹, Małgorzata Dudkiewicz¹, Dorota Mackiewicz¹, Maria Kowalczyk¹, Stanisław Cebrat¹, and Mirosław R. Dudek^{2*}

¹ Department of Genetics, Institute of Microbiology, University of Wrocław,
ul. Przybyszewskiego 63/77, PL-54148 Wrocław, Poland
{nowicka, pamac, malgosia, dorota, kowal, cebrat}@microb.uni.wroc.pl
<http://smORFland.microb.uni.wroc.pl>

² Institute of Physics, University of Zielona Góra, ul. A. Szafrana 4a,
PL-65516 Zielona Góra, Poland
mdudek@proton.if.uz.zgora.pl

Abstract. We have found that the effective survival time of amino acids in organisms follows a power law with respect to frequency of their occurrence in genes. We have used mutation data matrix PAM1 PET91 to calculate selection pressure on each kind of amino acid. The results have been compared to MPM1 matrix (Mutation Probability Matrix) representing the pure mutational pressure in the *Borrelia burgdorferi* genome. The results are universal in the sense that the survival time of amino acids calculated from the higher order PAM k matrices ($k > 1$) follows the same power law as in the case of PAM1 matrices.

1 Introduction

Determining the evolutionary distances between two protein sequences requires the knowledge of the substitution rates of amino-acids. It is generally accepted that the more substitutions are necessary to change one sequence into another, the more unrelated they are and the larger their distance to the common ancestor. The most widely used method for the calculation of distances between sequences is based on the mutation data matrix, M_{ij} , published by Dayhoff et al. [1], where i, j represent amino acids, and an element M_{ij} of the matrix gives the probability that the amino acid in column j will be replaced by the amino acid in row i after a given evolutionary time interval. The interval corresponding to 1 percent of substitutions between two compared sequences is called one PAM (Percent of Accepted Mutations), and the corresponding matrix is denoted as PAM1 matrix. There is assumed a Markov model of sequence evolution and a simple power M^k of the PAM1 matrix (multiplied by itself k times) denotes a matrix, PAM k , that gives the amino acid substitution probability after k PAMs. Today, a much more accurate PAM matrix is available, generated from 16130

* corresponding author

protein sequences, published by Jones et al. [2]. The large number of compared genes guarantees that the matrix has negligible statistical errors and it can be considered to be the reference matrix during the calculations of the phylogenetic distances. The matrix is also known as PET91 matrix.

Recently, by comparing intergenic sequences being remnants of coding sequences with homologous sequences of genes, we have constructed an empirical table of the nucleotide substitution rates in the case of the leading DNA strand of the *B. burgdorferi* genome [3],[4],[5]. We have found that substitution rates, which determine the evolutionary turnover time of a given kind of nucleotide in third codon positions of coding sequences, are highly correlated with the frequency of the occurrence of that nucleotide in the sequences. There is a compositional bias produced by replication process, introducing long-range correlation among nucleotides in the third positions in codons, which is very similar to the bias seen in the intergenic sequences [6].

We have used the empirical table of nucleotide substitution rates to simulate mutational pressure on the genes lying on the leading DNA strand of the *B. burgdorferi* genome and we have constructed MPM1 matrix (Mutation Probability Matrix) for amino acid substitutions in the evolving genes. Thus the resulting table represents the percent of amino acid substitutions introduced by mutational pressure and not by selection. Next, we compared the survival times of the amino acids in the case without any selection with the effective survival times of the amino acids, counted with the help of the PAM1 PET91 matrix.

2 Mutation Table for Nucleotides

DNA sequence of the *B. burgdorferi* genome was downloaded from the website www.ncbi.nlm.nih.gov. The empirical mutation table for nucleotides in third positions in codons, which we used in the paper, is the following [3],[4]:

$$M = \begin{pmatrix} 1 - uW_A & uW_{AT} & uW_{AG} & uW_{AC} \\ uW_{TA} & 1 - uW_T & uW_{TG} & uW_{TC} \\ uW_{GA} & uW_{GT} & 1 - uW_G & uW_{GC} \\ uW_{CA} & uW_{CT} & uW_{CG} & 1 - uW_C \end{pmatrix} \quad (1)$$

where ¹

$$\begin{aligned} W_{GA} &= 0.0667 & W_{GT} &= 0.0347 & W_{GC} &= 0.0470 & W_{AG} &= 0.1637 \\ W_{AT} &= 0.0655 & W_{AC} &= 0.0702 & W_{TG} &= 0.1157 & W_{TA} &= 0.1027 \\ W_{TC} &= 0.2613 & W_{CG} &= 0.0147 & W_{CA} &= 0.0228 & W_{CT} &= 0.0350 \end{aligned} \quad (2)$$

and the elements of the matrix give the probability that nucleotide in column j will mutate to the nucleotide in row i during one replication cycle. The symbols W_{ij} represent relative substitution probability of nucleotide j by nucleotide i , and u represents mutation rate. The symbols W_j in the diagonal represent relative substitution probability of nucleotide j :

¹ The transpose matrix convention has been chosen in [3].

$$W_j = \sum_{i \neq j} W_{ij}, \quad (3)$$

and $W_A + W_T + W_G + W_C = 1$.

The expression for the mean survival time of the nucleotide j depends on W_j as follows (derivation can be found in [3])

$$\tau_j = -\frac{1}{\ln(1 - u W_j)} \approx \frac{1}{u W_j}. \quad (4)$$

The above approximated formula is true for small values of the mutation rate u .

In papers [3],[4],[5], we concluded that in a natural genome the frequency of occurrence f_j of the nucleotides, in the third position in codons, is linearly related to the respective mean survival time τ_j ,

$$f_j = m_0 \tau_j + c_0, \quad (5)$$

with the same coefficients, m_0 and c_0 , for each nucleotide. The Kimura's neutral theory [7] of evolution assumes the constancy of the evolution rate, where the mutations are random events, much the same as the random decay events of the radioactive decay. However, the linear law in (5) is not contrary to the Kimura's theory. Still, the mutations represent random decay events but they are correlated with the DNA composition.

3 Mutational Pressure MPM1 Matrix Construction for Amino Acids

We used computer random number generator to generate random deviates with a distribution defined by the elements of the mutation matrix in (1). With the help of the random deviates we were mutating nucleotide sequences of 564 genes from the leading DNA strand of the *B. burdorferi* genome. The applied value of the mutation pressure was $u = 0.01$.

For each gene, considered to be an ancestral one in this simulated evolution, we prepared 10^5 pairs of homologous sequences, which diverged from this gene. The gene evolution was stopped when the number of substitutions between the homologous protein sequences reached 1%. All the sequences were translated into amino acids and we constructed a mutation probability matrix MPM1 according to the procedure of Dayhoff et al.[1] and Jones et al. [2]. The resulting mutation table, with substitution probabilities M_{ij} , the amino acid mutability m_j , and the fraction f_j of amino-acid in the compared sequences have been presented, respectively, in Table 1 and Table 2.

The elements M_{ij} of the MPM1 matrix in Table 1 have been scaled with the parameter λ , which related them to the evolutionary distance of one percent of substitutions and it is equal to 0.00009731 in our simulations. We introduced the parameter λ following the equation (3) in the paper by Jones et al.[2]). Therefore,

Table 1. Mutation Probability Matrix for an evolutionary distance of 1 PAM (splitted into two parts). Values of the matrix elements are scaled by a factor of 10^5 and rounded to an integer. The symbols in the first row and the first column represent amino-acids and numbers following colons -number of codons representing a given amino-acid in the universal genetic code.

	A:4	R:6	N:2	D:2	C:2	Q:2	E:2	G:4	H:2	I:3
A:4	99027	0	0	79	1	0	57	78	0	1
R:6	0	98773	1	0	251	139	1	218	175	51
N:2	0	2	98926	336	4	1	1	1	308	162
D:2	101	0	338	98936	4	0	297	215	184	0
C:2	0	50	0	0	97442	0	0	87	1	0
Q:2	0	76	0	0	0	99244	41	0	500	0
E:2	83	2	1	340	0	151	99125	247	1	0
G:4	92	357	1	199	721	0	200	99103	0	0
H:2	0	56	56	33	2	295	0	0	98315	0
I:3	1	153	277	1	3	0	0	1	1	99026
L:6	1	90	0	0	4	198	0	0	459	150
K:2	0	481	452	1	0	279	354	1	1	122
M:1	0	69	0	0	0	0	0	0	0	137
F:2	1	0	1	0	697	0	0	0	2	165
P:4	55	52	0	0	1	82	0	0	180	0
S:6	218	324	215	1	1070	0	0	161	2	79
T:4	297	46	71	0	1	0	0	0	0	163
W:1	0	63	0	0	190	0	0	41	0	0
Y:2	0	0	235	198	868	1	1	0	591	0
V:4	435	1	1	213	2	0	169	235	1	225

	L:6	K:2	M:1	F:2	P:4	S:6	T:4	W:1	Y:2	V:4
A:4	0	0	0	0	107	129	333	0	0	279
R:6	29	177	116	0	73	140	38	653	0	0
N:2	0	292	1	1	0	163	102	0	330	1
D:2	0	1	0	0	0	1	0	0	279	176
C:2	0	0	0	62	0	91	0	389	137	0
Q:2	34	56	0	0	63	0	0	1	1	0
E:2	0	263	1	0	0	0	0	3	1	159
G:4	0	1	0	0	0	113	0	686	1	179
H:2	47	0	0	0	81	0	0	0	151	0
I:3	143	135	693	223	1	102	401	0	1	317
L:6	99268	0	283	515	461	130	1	593	1	150
K:2	0	99246	287	0	0	1	107	3	1	1
M:1	53	63	98686	0	0	0	42	2	0	46
F:2	363	0	1	98928	2	211	1	2	303	149
P:4	104	0	0	1	98949	148	85	0	0	0
S:6	96	1	1	221	485	98951	286	91	117	1
T:4	0	48	86	0	146	151	98725	0	0	1
W:1	18	0	0	0	0	4	0	98760	0	0
Y:2	1	1	0	171	0	63	0	2	99034	0
V:4	102	1	166	143	1	1	2	2	1	98798

the diagonal elements of the MPM1 table have been defined with the help of the following formula,

$$M_{jj} = 1 - \lambda m_j, \quad (6)$$

and the value of the parameter λ has been calculated from the condition that, in the case of 1% of amino acid substitutions, the total fraction of the unchanged amino acids is equal to

$$\sum_{j=1}^{20} f_j M_{jj} = 0.99. \quad (7)$$

Table 2. Relative mutabilities and fractions of 20 amino acids in the compared sequences. We used the convention that mutabilities are relative to alanine and it is arbitrarily assigned a value of 100.

amino acid	relative mutability (m_j)	fraction (f_j)
A	100.00	0.0449
R	126.09	0.0369
N	110.42	0.0671
D	109.29	0.0579
C	262.91	0.0073
Q	77.67	0.0206
E	89.88	0.0644
G	92.21	0.0582
H	173.18	0.0114
I	100.12	0.0964
L	75.23	0.1060
K	77.51	0.0930
M	135.01	0.0184
F	110.12	0.0690
P	108.02	0.0238
S	107.78	0.0796
T	131.05	0.0333
W	127.44	0.0047
Y	99.25	0.0427
V	123.56	0.0645

4 Discussion of results

The major qualitative difference between the MPM matrix introduced in the previous section and the PAM1 PET91 matrix published in the paper by Jones et al.[2] is that the first one is a result of pure mutational pressure whereas the

second one is a result of both mutational and selection pressures. Thus, we have two evolutionary mechanisms responsible for the resulting PAM matrices.

With the help of formula (4) (extended to amino acids) we have calculated effective survival times of amino acids in the case of the MPM1 matrix (Table 1) and the mutational/selectional PAM1 PET91 matrix ([2]). The value of the parameter λ is a counterpart of u in (4). In Fig.1, we presented the relation between the calculated survival time of amino acids and their fractions in the *B. burgdorferi* proteins, in the pairs of diverged genes, in a log-log scale. One can observe that the data are highly correlated and in both cases the dependence of the mean survival time of amino acid on the fraction of the amino acid represents a power law:

$$\tau_j \sim F_j^\alpha \quad (8)$$

with a negative value of $\alpha \approx -1.3$ in the case of selection and a positive value of $\alpha \approx 0.3$ in the case of mutation pressure on the leading DNA strand of the *B. burgdorferi* genome. The value of α for the analogous mutational PAM1 matrix calculated in the case of the lagging DNA strand of the *B. burgdorferi* genome is about twice as small. It is worth to underline that the slopes α are the same for the matrices PAM k with high values of k , and thus, they are universal with respect to evolution.




Fig. 1. Relation between survival time of amino-acids and their fractions in compared pairs of homologous genes in the case with selection and in the case without selection. Selection data for amino acids have been taken from PAM1 PET91 matrix whereas the data in the case without selection have been simulated using experimentally found mutational pressure of the *B. burgdorferi* genome.




Fig. 2. Gene length versus gene rank, where each gene has assigned a rank with respect to fraction of tryptophane (left graph) and with respect to leucine (right graph). In each graph there are two plots: the dots represent gene's size vs. rank, whereas the second plot represent the fraction vs. rank.

In Fig.1 we may observe a kind of evolutionary scissors acting on amino acids. Once the less frequent amino acids, like *W* (tryptophane), *C* (cysteine), have much shorter turn over time compared with other amino acids (as can be seen from the lower line) the selection pressure (upper line) counteracts with the effect. On the other hand, the most mutable amino acids, like *L* (leucine) or *I* (isoleucine), which are very frequent in genes, seem to be much weakly influenced by selection.

In genes the fraction of the amino acids most protected by selection strongly depends on gene size, it is diminishing when gene's size is increasing. The effect weakens if we go into right the direction of the evolutionary scissors in Fig.1.

To show this, we ordered all 564 genes under consideration with respect to fraction of an examined amino acid and the genes have been assigned a rank number. Next, we plotted the dependence of both the gene size on the rank and the dependence of the amino acid fraction in the gene on the rank. The resulting plots in Figs.2 correspond to two evolutionary extreme cases, representing tryptophan and leucine. It is evident that in the case of tryptophan the fraction of that amino acid in genes is anti-correlated with the gene's size (notice, that about 1/3 genes do not possess tryptophan). In the case of leucine there is a crossover and the effect of selection is evident only for the genes which have more than 10% of that amino acid. When the fraction of leucine is less than 10% there is even a reverse effect, i.e., the increasing fraction is correlated with the increasing gene's size. If we look at the evolutionary scissors in Fig.1, we can see that in the case of leucine the survival time, which originates from pure

mutational pressure is longer than its selectional counterpart. Recently, there has appeared a paper by Xia and Li [8] discussing which amino acid properties (like polarity, isoelectric point, volume etc.) affect protein evolution. Thus, there is a possibility to relate these properties to our discussion of the evolutionary scissors.

5 Conclusions

We have shown that the amino acids which experience the highest selectional pressure have the shortest turn over time with respect to mutation pressure. The fraction of these amino acids in genes depends on the gene's size. Much different is the selectional role of the amino acids, like leucine, from the right hand side of the selectional scissors. Although they have long turn over time with respect to the mutational pressure, their fraction cannot be too high. This could be considered as an effect of optimisation of the genetic information on coding with processes of mutagenesis and phenotype selection for protein functions.

Acknowledgements

The work was supported by the grant number 1016/S/IMi/02.

References

1. Dayhoff, M.O., Schwartz, R.M., and Orcutt, B.C.: A Model of Evolutionary Change in Proteins. In: Atlas of Protein Sequence and Structure , Vol. 5 Suppl. 3 (1978) 345–352
2. Jones, D.T., Taylor, W.R., and Thornton, J.M.: The rapid generation of mutation data matrices from protein sequences. In: CABIOS, Vol. 8 no. 3, (1992) 275–282
3. Kowalczyk, M., Mackiewicz, P., Szczepanik, D., Nowicka, A., Dudkiewicz, M., Dudek, M.R., and Cebrat, S.: Multiple base substitution corrections in DNA sequence evolution. *Int. J. Mod. Phys. C* **12** (2001) 1043–1053
4. Mackiewicz, P., Kowalczyk, M., Mackiewicz, D., Nowicka, A., Dudkiewicz, M., Łaskiewicz, A., Dudek, M.R., Cebrat, S.: Replication associated mutational pressure generating long-range correlation in DNA. *Physica A* **314** (2002) 646–654
5. Kowalczyk, M., Mackiewicz, P., Mackiewicz, D., Nowicka, A., Dudkiewicz, M., Dudek, M.R., and Cebrat, S: High correlation between the turnover of nucleotides under mutational pressure and the DNA composition. *BMC Evolutionary Biology* **1** (2001) (1):13
6. Cebrat, S., Dudek, M.R., Gierlik, A., Kowalczyk, M., Mackiewicz, P.: Effect of replication on the third base of codons. *Physica A* **265** (1999) 78–84
7. Kimura, M.: The Neutral Theory of Molecular Evolution, Cambridge University Press, Cambridge (1983)
8. Xia, X., Li, W.-H.: What Amino Acid Properties Affect Protein Evolution. *J. Mol. Evol.* **47** (1998) 557–564